

Durham Research Online

Deposited in DRO:

23 April 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Abedi, Sepideh and Astarai, Fatemeh Razi and Ghobadian, Barat and Tavakoli, Omid and Jalili, Hassan and Chivasa, Stephen and Chris, Greenwell H. (2019) 'Bioenergy production using *Trichormus variabilis* - a review.', *Biofuels, bioproducts biorefining.*, 13 (5). pp. 1365-1382.

Further information on publisher's website:

<https://doi.org/10.1002/bbb.2023>

Publisher's copyright statement:

This is the accepted version of the following article: Abedi, Sepideh, Astarai, Fatemeh Razi, Ghobadian, Barat, Tavakoli, Omid, Jalili, Hassan, Chivasa, Stephen Greenwell H. Chris (2019). Bioenergy Production Using *Trichormus variabilis* - A review. *Biofuels, Bioproducts Biorefining* 13(5): 1365-1382, which has been published in final form at <https://doi.org/10.1002/bbb.2023>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Bioenergy Production Using *Trichormus variabilis* - A review

Sepideh Abedi^{1,6}, Fatemeh Razi Astarai^{1*}, Barat Ghobadian², Omid Tavakoli³, Hassan Jalili⁴, Stephen Chivasa^{5*}, H. Chris Greenwell⁶

¹Department of Renewable Energies and Environmental Engineering, Faculty of New Sciences and Technologies, University of Tehran, Tehran, Iran.

²Department of Biosystems Engineering, Tarbiat Modares University, Tehran, Iran.

³School of Chemical Engineering, College of Engineering, University of Tehran, Tehran, Iran.

⁴Department of Life Science Engineering, Faculty of New Sciences and Technologies, University of Tehran, Tehran, Iran.

⁵Department of Biosciences, Durham University, Durham, DH1 3LE, United Kingdom.

⁶Department of Earth Sciences, Durham University, Durham, DH1 3LE, United Kingdom.

Running Head: Bioenergy from *Trichormus variabilis*

***Corresponding authors:**

Fatemeh Razi Astarai: +98-912-297-7929; Fax: +98-21-884-97324

E-mail: razias_m@ut.ac.ir

Stephen Chivasa: Tel: +44-191-334-1275; Fax: +44-191-334-1201

E-mail: stephen.chivasa@durham.ac.uk

Abstract

Fossil fuel processing and consumption contaminates air, soil, and water resources via release of hazardous chemicals. To protect the environment, harnessing renewable energy resources and development of sustainable technologies are prime targets of research and increased investment. Use of bio-based feedstocks in energy production inherently provides a valuable pollution-curbing pathway with sustainability credentials, especially when wastewater is used to provide the nutrient requirements. The filamentous cyanobacterium, *Trichormus variabilis*, has attracted substantial attention from researchers due to its potential for dual industrial functions in bioenergy production and bioremediation. This species can efficiently use the power of sunlight energy to fix atmospheric CO₂ for generating valuable chemical compounds, such as carbohydrates and fatty acids, which can be converted to biofuels. Because it grows in nutrient-rich wastewater (industrial effluent), it can serve as a bio-absorbent and replace costly chemical catalysts and nano-materials classically used for removal of nutrients and metals. However, no recent review has presented potential for state-of-the-art *T. variabilis*-driven phycoremediation-bioenergy production systems. Therefore, in order to present possible routes from phycoremediation to energy production as a strategy for developing the industrial application of *T. variabilis*, we present this review to bring important research results on this species together and highlight major related challenges and opportunities. The current status of applying algae in bioremediation and production of liquid and gaseous fuels utilizing wildtype and mutants of *T. variabilis* is explored. Finally, key points underlying potential for future research on optimization of robust technologies for supplying sustainable bioenergy using this organism are presented.

Key words: Bioenergy, Hydrogen, Bioremediation, *Trichormus variabilis*.

1 1 Introduction

2 Global economic growth over the last two centuries has largely been driven by
3 increased use of fossil fuels and was accompanied by significant environmental
4 pollution, particularly increased CO₂ emissions. Total world energy consumption is
5 forecast to increase by 1.4% per annum up to 2040¹, exacerbating negative
6 environmental impacts and increasing pressure on finite fossil fuel resources.
7 Additionally, both economic growth and rises in the global population are reciprocated
8 by increased demand for freshwater required for industrial processes, agriculture, and
9 domestic use. Therefore, recycling wastewater becomes an imperative for water
10 conservation. There exist a number of routes to transform the environmental challenges
11 posed by wastewater into efficient, economically viable methods for both energy
12 production and water recycling. Phycoremediation has been used since circa 1963 as a
13 cost-effective nutrient removal method using microalgae to treat municipal wastewater².
14 In addition to CO₂ capture, phycoremediation has several other benefits, including
15 removal, transformation, or degradation of nutrients, organic matter, acids, metals and
16 xenobiotic compounds, and the use of algae as an environmental monitoring system for
17 detection of toxic materials³ and production of high-value metabolites⁴. Furthermore,
18 massive biomass grown in wastewater streams can be used effectively as a promising
19 source of bio-fuels such as bioethanol, biodiesel, bio-hydrogen or biogas, which are
20 generally known as the most carbon neutral liquid and gaseous fuels, depending on their
21 generation technology.

22 Previous reviews have collated research on phycoremediation⁵, bioenergy
23 production^{6, 7}, biorefining⁸ and algal photobioreactors (PBRs)⁷. However, to our
24 knowledge, none of these reviews have been dedicated to a specific strain of algae.
25 Among many microalgae species, *Trichormus variabilis* has been studied extensively
26 owing to its dual benefits: showing high potential in industrial and municipal
27 wastewater treatment⁹⁻¹¹ and also its capability for bio-hydrogen production^{12, 13}. The
28 worldwide distribution of *T. variabilis*, high auto-flocculation capacity³, tolerance of a
29 wide temperature range, and bio-fertilizer potential¹⁴ strongly affirm that the study of this
30 species needs to be expanded in the future. Here, we present a comprehensive review,
31 focused on *T. variabilis*, and its principal enzymatic functions which make it suitable
32 for applications in biohydrogen production. Subsequently, we present the physico-
33 chemical parameters affecting the morphology, physiology and growth of *T. variabilis*
34 and evaluate how these factors might impinge on the hydrogen production process.
35 Finally challenges and opportunities in developing a route to industrial application of *T.*
36 *variabilis* as a green chemistry workhorse in effluent treatment and production of
37 sustainable bioenergy are discussed.

38 2 Methodology

39 Primary and secondary literature and data used in synthesis of this review were
40 obtained via searching Web of Science (<https://clarivate.com/products/web-of-science/>)
41 and PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/>) databases and several sources of
42 grey literature using keywords related to *T. variabilis* in terms of various types of
43 bioenergy and nutrient removal potential. Synonyms of *T. variabilis*, such as *Anabaena*
44 *variabilis*, were also used as keywords given that authors tend to use different names for
45 this organism. Information was gathered in a two-phased process. In the first phase,
46 inclusion criteria were peer-reviewed academic research and published industrial pilot
47 studies or full-scale production systems. Because of paucity of published data specific
48

to *T. variabilis* no exclusion criteria were applied, except the exclusion of self-published, non-peer-reviewed material. In order to capture important data from grey literature, such as patents and technical reports, the second phase reversed the inclusion and exclusion criteria of the first phase. The taxonomic classification of *Trichormus variabilis* was extracted from data presented in major biological, botanical and algal databanks including AlgaeBase¹⁵, World Register of Marine Species (WoRMS ID: 146661)¹⁶ and National Center for Biotechnology Information (NCBI taxonomy ID: 240292, <https://www.ncbi.nlm.nih.gov/>). Genes encoding hydrogen biosynthetic enzymes (described in Section 6.3) were identified via literature and Genome Database Resources of CyanoBase in Kazusa DNA Research Institute (<http://genome.microbedb.jp/cyanobase/>). Vector graphics were drawn by Inkscape software v0.92¹⁷. To provide images of *T. variabilis* cells, algal samples were cultivated and processed for transmission electron microscopy (TEM) as described previously¹⁸. All the tables in this review were constructed from the primary results reported in different studies cited within each tables. Figure 1 presents a workflow chart illustrating the search strategy used.

Figure1. Flowchart of the research strategy. To develop each section of this literature review, a question was formulated and related keywords generated for interrogation of the databases. The returned information hit list was divided into three broad categories of algal cultivation for biomass production, algal cultivation for bioenergy production, and algal cultivation for phycoremediation. Progress of innovation across these were critically analysed by comparative analysis and current challenges and opportunities identified. Areas of future research to ensure current investments lead to commercial feasibility were finally identified and discussed.

3 *Trichormus variabilis*

Trichormus variabilis (syn *Anabaena variabilis* ATCC 29413) is a filamentous cyanobacterium from Nostocoidae subfamily^{19, 20} found in both freshwater and within soil habitats. The recorded history of study of this species begins nearly two centuries ago, when it was collected for the first time in 1839 from puddles and drains in Hooksiel and Jadebusen, located in Wangerooge Island in Lower Saxony of Germany by Kutzing²¹. Since then, its widespread distribution has been reported across almost all continents, being a native species in many countries, from Asia to America, Africa and Australia^{22, 23}. The taxonomy of this species has faced repeated revision owing to the formation of two main cell types (akinetes and heterocysts), including differentiation from the genus of *Nostoc*²⁴, *Wolleea*²⁵ and *Anabaena*^{19, 20, 24, 25} on the basis of akinete formation. According to the last rectified taxonomy (Fig. 2), it has been transferred from *Anabaena* to the genus *Trichormus*, with the former scientific name “*Anabaena variabilis*” proposed by Bornet and Flahault in 1886 being replaced with “*Trichormus variabilis*” as proposed by Komarek and Anagnostidis in 1989²⁵⁻²⁷.

Figure 2. Taxonomy of *Trichormus variabilis*.

3.1 Morphology and Physiology

The filaments of *T. variabilis* are morphologically variable^{19, 24}. Filaments mainly contain chains of barrel-shaped gram-negative vegetative cells (Fig. 3a) as a site for

photosynthesis with conical terminal cells (Fig. 3b) covered by a thin gelatinous polysaccharide sheath (Fig. 3c). The mucilaginous sheath encapsulating the filaments arises from polysaccharide secretions by the cells^{28, 29}. Vegetative cells have a primary role of harnessing solar energy for fixing carbon to drive biomass production. The growth in biomass can be coupled to wastewater treatment through removal of heavy metal ions and nutrients. The most important cell differentiation in *T. variabilis* is the formation of heterocysts, also known as heterocytes. When transferred to diazotrophic conditions, some vegetative cells differentiate into heterocysts as a site for nitrogen fixation²⁵. The spherical, ellipsoidal or cylindrical shape²⁴, the large size²⁵, distinctive polar bodies³⁰, and hollow appearance, make heterocysts readily recognisable among other cell types (Fig. 3d). Even though heterocysts represent only 5-10% of cells along the filaments³¹, phylogenetic studies showed that about 45% of the genome is expressed in these cells³². Cell differentiation is underpinned by changes in gene expression, which trigger biochemical and structural transformations, such as formation of thick multi-membranous glycolipid walls, arrest of photosynthesis and biosynthesis of the nitrogenase enzyme complex³³. These modifications prepare heterocyst cells for the vital role of fixing atmospheric nitrogen with hydrogen gas as a by-product of the reactions. Because nitrogenase enzymes are sensitive to oxygen, heterocysts bereft of oxygenic photosynthesis provide an anaerobic environment to stabilise the N-fixation system in an otherwise aerobic organism. Heterocysts strongly attach to vegetative cells via cyanophycin polar nodule³⁰ (Fig. 3d).

In response to nutrient limitation or environmental stress such as prolonged darkness, dryness, freezing, and nutrient depletion^{30, 34, 35}, specialised akinete cells develop as resilient archetypal spores to ensure survival of the organism in harsh environmental conditions. Akinetes are oval to barrel-shaped cells with granules filled with glycogen and cyanophycin³⁶, which can easily be distinguished from vegetative cells by their larger size and thicker envelope (Fig. 3e). Their thick outer layer contains glycolipid and polysaccharides, making them more resistant to stressful conditions in comparison to vegetative cells and heterocysts²⁸. Akinetes germinate and develop into new filaments when favourable growth conditions are restored. On germination, an akinete develops into a short filament which is structurally distinguishable from trichorms in having 5 to 15 vegetative cells and lacking heterocysts (Fig. 3f). These short filaments, known as hormogonia, later develop into mature trichorms possessing heterocysts. - Therefore, the diverse cell functions provide an attractive system for commercial development of an efficient low-cost biomass production during wastewater treatment process, which could be coupled to industrial bioenergy production. The morphological appearance of various cell types of *T. variabilis* is illustrated schematically in Fig. 3, with Fig. 4 showing actual images of the cells using transmission electron microscopy (TEM).

Figure 3. Schematic depiction of different cell types of *T. variabilis*. (a) Vegetative cells; (b) Terminal cell; (c) polysaccharide mucilaginous sheath; (d) Heterocysts, which posses: 1. Fibrous layer, 2. Laminated layers (internal layers), 3. Plasmodesmata, 4. Polar nodule, and 5. Thylakoids; (e) Akinete, which has: 6. sheath, 7. additional wall, 8. Cyanophycin granule, 9. Inner layer, and 10. Dense fibrillar layer; (f) Hormogonium; (g) Outer membrane; (h) Polyphosphate granules.

Figure 4. Microscopic images of *T. variabilis* cells. (a) Low power light microscope image of filaments in nitrogen-depleted growth medium showing vegetative

cells, heterocysts, and akinete. **(b)** Transmission electron micrograph of filaments encased by a gelatinous polysaccharide sheath (PC) with terminal cells (TC) at the end of the trichorms. **(c)** Transmission electron micrograph showing a detached mature vegetative cell showing peptidoglycan layer (PG), carboxysome (CS), polyphosphate granule (PS), cyanophycin granule (CP), thylakoid and attached phycobilisomes (Tl). Scale bars: **(a)** 50 μm ; **(b)** 2 μm ; **(c)** 500 nm.

4 Physico-chemical factors affect *T. variabilis* biomass production

Knowing key physico-chemical factors required for promoting high rates of biomass production is necessary for developing industrial-scale bioreactors for algal cultivation. Zho³⁷ summarized general microalgae cultivation strategies, abiotic, biotic and operational factors affecting algal growth. *T. variabilis* is capable of growing phototrophically, heterotrophically/chemotrophically in the dark with exogenous sugar, and also mixotrophically in light with exogenous sugar^{31, 38, 39}. This variation in growth strategy makes it a potentially favourable microorganism for use in different light- and dark-driven bioenergy production processes. *T. variabilis* is a mild thermophilic cyanobacterium, which has photosynthetic activity in a broad temperature range (10- 35 °C) and light intensity (42- 562 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$)⁴⁰. It was demonstrated that the growth yield declined (0.1 g dry mass.g⁻¹ CO₂) by increasing the initial CO₂ concentration from 4 to 18%⁴¹. Chemical nutrients, essential salts and metals directly influence different metabolic modes, which consequently affects bioenergy yield and remediation potential. The main growth media used for *T. variabilis* include blue-green (BG)⁴², Allen and Arnon (AA)⁴³, 2-(N-Morpholino) ethane sulfonic acid-Volvox (MES)⁴⁴, Marin (MN)⁴⁵, ammonium mineral salt (AMS-1)⁴⁶ and Fogg's Medium⁴⁷. Among these, BG and AA are the most commonly used media, specifically when *T. variabilis* is used as a source of bio-hydrogen. Higher biomass production was reported when grown in BG₁₁ enriched with a nitrogen source⁴⁸. In general, BG and AA media are based on molybdenum (Mo) and vanadium (V), respectively. These two metals are required for activation of two key enzymes (described in section 5.3), which are responsible for hydrogen generation. Investigating the effects of replacing Mo with V demonstrated that replacement of Mo with V leads to decreases in the growth yield⁴³ while it improves hydrogen generation yield⁴⁸ (section 5.4).

Experiments establishing *T. variabilis* growth profiles have classically used, as proxies for biomass accumulation, parameters such as photosynthetic potential/capacity⁴⁹ and cell density at specified wavelengths⁵⁰. These have been applied in studying the effects of altering various physicochemical factors such as light and nutrient supply, on *T. variabilis* phototrophic^{51, 52} or heterotrophic^{39, 53} growth. While a large number of studies have investigated *T. variabilis* growth rate as a function of time, cell density or photosynthesis components^{54, 55}, only a few have directly surveyed the interrelation between multiple parameters. Such data can be used for modelling growth kinetics. In prior work, a kinetic model for growth of *T. variabilis* was developed based on the Monod model as a function of CO₂ and light intensity⁵⁶. Predictive models are increasingly important as algal biomass generated from wastewater treatment process is being used for production of biofuels when scaling-up from laboratory-scale to industrial-scale cultures. Therefore, the model should capture key parameters affecting growth rate so that bioreactor designs are optimised for maximal productivity.

Additionally, this species has variable morphology based on the type of mechanical stimulation. It normally grows as filaments that attach longitudinally to neighbouring

filaments to form thalloid structures, but when it is grown with orbital shaking, thalli formation is prevented, with a homogenous suspension of short filaments emerging⁴³. High tendency of filaments for sticking to the walls blocks light penetration, which causes problems when cultivating this species in photobioreactors. The membranous growth form may provide an opportunity in flocculation of biomass in open raceway ponds. This potential, which resolves one of the most important challenges of industrial algal harvesting technologies in biofuel production, requires further investigation.

5 Biomass production via phycoremediation

Providing sufficient water and nutrients, particularly low-cost nitrogen, is a challenge for large-scale microalgae cultivation for bioenergy⁷. However, nutrient-rich wastewater streams can provide most of the nutrients required for algal growth. In addition to utilising nitrate and phosphate, microalgae have the capacity for effective removal of heavy metals. Incorporation of metals into the biomass may pose downstream risks during future processing or utilisation of derivatives, depending on the uses of the biomass. Owing to their rapid growth rates and potential to remove heavy metals, various microorganisms have been used in phycoremediation of municipal⁵⁷, industrial⁵⁸ or agricultural⁵⁹ wastewaters. These microorganisms include (i) green algae; *Chlamydomonas reinhardtii*⁶⁰, *Chlorella sp.*⁶¹, *Nannochloropsis oculata*⁶², *Scenedesmus dimorphus*⁵⁹, (ii) blue-green algae; including *Synechocystis salina*⁶³, *Spirulina platensis*⁶⁴, *Phormidium bohneri*⁶⁵, *Tolypothrix ceytonica*⁹, (iii) brown algae, including *Padina sp.*⁶⁶, *Laminaria hyperborean*⁶⁷, and (iv) red algae such as *Porphyridium cruentum*⁶⁸. Among the cyanobacteria, *T. variabilis* has received considerable attention for use in bioremediation. Attraction to this species has partly been the potential for coupling wastewater treatment with production of bioenergy. The growth and remediation performance of *T. variabilis* in treating water contaminated with ammonium⁶⁹ and biosorption of heavy metals, such as Cr, Cd, Ni and Pb⁷⁰ has been demonstrated. Therefore, *T. variabilis* is already noted as an efficient microalgae for actual and potential treatment of mixed industrial and municipal wastewater⁹. Table 1 presents a synopsis of key studies on phycoremediation using *T. variabilis* and the maximum removal efficiency reported for each process.

A study on phenol compounds remediation illustrated notable nitrophenol compound removal efficiency from an artificial supplemented wastewater using this species⁷¹. However, since polyphenolic compounds produced by *Myriophyllum verticillatum* cause growth inhibition of *T. variabilis* cells⁷², the feasibility and efficiency of *T. variabilis* remedial action on phenolic contaminants will require further research. It was also reported that nanomaterials, such as nano-titanium dioxide (nTiO₂), have negative effect on the intracellular structure of *T. variabilis*⁷³⁻⁷⁵. Furthermore, *T. variabilis* growth is impaired by some allelochemicals such as Harmane (1-methyl- β -carboline)⁷⁶ and FischerellinB⁷⁷ causing cell lysis and photosynthesis inhibition respectively. Microcystins have intense allelopathic inhibition on *T. variabilis* growth⁷⁸ and differentiation of heterocysts and akinetes⁷⁹. The presence of such pollutants in wastewater streams affects nutrient removal efficiency by disrupting metabolic processes and consequently, disrupts bioenergy production pathways. Therefore, more research on detection of growth inhibiting pollutants, their effects, and ways to deal with them is required. Exposing *T. variabilis* to pollutants can gradually change gene expression and acclimate the organism for growth in previously toxic compounds. For example, it was demonstrated that repeated cultivation of this species in a medium with

high concentration of $\text{Cu}(\text{NO}_3)_2$ developed a Cu-resistant strain with broad resistance to Cd, Zn and Ni⁴⁹. Thus, genomics and proteomics research could provide a clear understanding of the exact metabolic changes underpinning this adaptive change.

6 Bioenergy production using *T. variabilis*

Several studies focused on integration of wastewater treatment and bio-fuel production using microalgal systems⁸⁰⁻⁸³. After nutrient removal, the biomass produced can effectively be utilized as a resource for cost-effective bioenergy production. We review the application of *T. variabilis* in bioenergy production from the perspective of liquid (biodiesel, bio-ethanol) and gaseous (bio-hydrogen, bio-methane) bio-fuels. Applying microalgae systems for bioenergy production has several environmental advantages, such as reduction of pollutant emissions and soil erosion, and economic benefits such as tax credits and grants without competing with agricultural produce for freshwater and arable land^{84, 85}. In addition to these general advantages, *T. variabilis* has enhanced capacity for auto-flocculation³ which is one of the most important characteristics to decrease energy consumption during algae biomass harvesting. Also, nutrient removal and CO_2 bio-fixation potential together with interesting cell differentiations in forming heterocysts and akinetes, which provide suitable opportunities for N-fixation and growth in challenging environmental conditions, respectively, and availability of processed biomass as bio-fertilizer¹⁴, suggest this species may present a viable solution for algae bioenergy production with valuable by-products.

6.1 Liquid bio-fuels: biodiesel and bioethanol

Ethical concerns about diversion of crops from the human food chain to fuel production have spurred the search for inedible crop oils and waste fats for use as a bio-fuel feedstock. Inedible oilseeds, such as *Jatropha*, *Castor*, and other indigenous crops^{86, 87} which can grow on marginal land have emerged as potential candidates. However, the cost of land, labour and freshwater resources often make large-scale production uneconomical when set against the yields attained. Therefore, microalgae received greater attention and interest due to their capability to grow in wastewater. Since lipid content is the most important key factor in utilization of oleaginous microorganisms as a source of biodiesel production, considerable research focuses on increasing lipid yield. Although *T. variabilis* is not considered as an oleaginous species, previous reports indicate notable increases in lipid content after exposure to environmental stress or nutrient limitation. For example, it was demonstrated that lipid content of this alga in N-depleted Arnon medium at 30-35 °C under 12:12 h light/dark cycle was increased by ~ 2.5-fold (yield~10.5%)⁸⁸. Furthermore, it has been reported that ultrasound at 200 W after 5 minutes increased lipid content of *T. variabilis* grown in BG₁₁ medium at 25 °C under continuous illumination by 1.46-fold (yield~47%)⁸⁹. Cellular stress triggered by ultrasonic treatment was responsible for increased oil production while a longer duration of treatment damaged the cells. Such stress or nutrient limitation leading to increased lipid yield was also observed in other species, including *Nannochloropsis* sp.⁹⁰, *Chlorella* sp.⁹¹ and *Chlamydomonas reinhardtii*⁹². These useful results can be considered as a promising solution for promoting lipid accumulation in the cells and consequently enhance the biodiesel production yield. However this also requires accurate investigation on the profile of fatty acid methyl esters to match the biodiesel quality standards.

From a bio-ethanol perspective, *T. variabilis* has not been widely investigated. The main challenge in bio-ethanol production from this filamentous species is overcoming the polysaccharide loss during processing. Research has shown that drying the biomass using supercritical fluid followed by fermentation could increase the bio-ethanol yield (24.1%) about 2-fold comparing with using lyophilization process followed by fermentation (yield~13.6%)⁹³. A further study has confirmed that applying supercritical fluid pre-treatment efficiently enhances the amount of ethanol from 1.25 g.L⁻¹ to 2.28 g.L⁻¹⁹⁴. Evidently, more research is required to fully investigate the liquid bio-fuel production potential of *T. variabilis*, especially when it is grown in wastewater.

6.2 Gaseous bio-fuels: bio-methane and bio-hydrogen

Biogas is the oldest form of bioenergy generated by means of methanogenic bacteria under anaerobic conditions containing ~50-70% bio-methane, which can be used as a heat, power or transportation fuel⁹⁵. Several microalgae have been used to combine bio-methane production with biorefinery approaches, for example, *Chlamydomonas reinhardtii* and *Scenedesmus obliquus*⁹⁶. There are some reports of *T. variabilis* biomass conversion into biogas under high temperature using anaerobic digesters without⁹⁷ or in combination with immobilising technology to enhance gas production⁹⁸. Methane production yield was recorded at 450 mL.g⁻¹ biomass using immobilised methanogenic bacteria and *Rhodobacter capsulatus* on polymeric matrices in an anaerobic bioreactor⁹⁸. In another research cumulative methane yield (64%) was recorded at ~4 mmol.g⁻¹ biomass when anaerobic cellulolytic substrate were used together with methanogenic *Archaea* from genera of *Methanoculleus* and *Methanosarcina*⁹⁷. Biogas is generated via 4 successive stages: (i) hydrolysis of biopolymers to monomers, (ii) fermentation (acidogenesis) of amino acids and sugars to intermediary products, (iii) acetogenesis of intermediary products to acetate, CO₂ and hydrogen, and (iv) methanogenesis, which transforms acetate into methane. During anaerobic digestion, hydrolysis is known to be the rate limiting step which needs to be optimized via efficient pre-treatment technologies. Owing to the composition and structural features of microalgae resulting in changes in response to the different growth conditions⁹⁹, it is assumed that biogas production yield will be affected by *T. variabilis* grown in wastewaters prior to anaerobic digestion, through expression of various proteins. However, no research has yet demonstrated the effect of these generated macromolecules, which consequently impact bio-degradability of biomass during hydrolysis. Also, more investigation is required to be undertaken on the effect of co-digestion of *T. variabilis* and operational conditions such as hydraulic retention time and proportion of inoculum and substrates¹⁰⁰ as the main parameters known in bio-methane production. The results of previous studies indicated that variety and abundance of acetate oxidizing syntrophic bacteria significantly influence conversion efficiency of *T. variabilis* to bio-methane. Improving biogas conversion yield would be an interesting subject particularly in industrial wastewater treatment plants for coupling bioremediation with methane production.

The most common form of bioenergy generated from *T. variabilis* is hydrogen, a clean source of fuel that can be used either directly in internal combustion engines or in fuel cells to generate electricity¹⁰¹. The environmental benefits offered by renewable hydrogen are clear, particularly if the energy is being harnessed from microorganisms. Different species of algae¹⁰² have been identified as potential sources of bio-hydrogen. Basic research in this field started with investigating bacterial hydrogen production in

the 1920s and later in the 1940s, attention shifted to microalgal hydrogen production¹⁰³. Development in applied research started in the early 1970s¹⁰³. Across several species used as cyanobacterial sources of bio-hydrogen¹⁰⁴, *T. variabilis* has attracted much attention owing to its 2-fold benefits in bio-hydrogen production and its considerable potential in nutrient removal from wastewater⁹⁻¹¹. Table 2 presents hydrogen production yield of *T. variabilis* under different experimental conditions. Since this species has been widely investigated for hydrogen generation, here we focus on the biochemistry, cultivation methods, and utilization of wildtype and mutant strains for phycoremediation and/or fuel production.

6.3 Biochemistry and genetics of bio-hydrogen generation using *T. variabilis*

Algal hydrogen production occurs via light-dependent and light-independent reactions. Several articles have reviewed light- and dark-driven bio-hydrogen production¹⁰⁵⁻¹⁰⁸. The former includes bio-photolysis and photo-fermentation, which utilise sunlight, water, and CO₂ as energy, electron, and carbon sources, respectively. Light-independent hydrogen generation includes dark fermentation, which uses carbohydrate substrates, such as found in organic waste materials. Light-dependent processes are mainly driven by photosynthesis¹⁰⁹. Hydrogen can be generated aerobically as a product of oxygenic photosynthesis or under anaerobic conditions, as a by-product of the conversion of organic substrates to acidogenic materials^{106, 110}. Here, we have reviewed H₂ production methods using *T. variabilis* with specific focus on the biochemistry of enzyme systems.

The complete genome of *T. variabilis* ATCC 29413 was sequenced by the department of energy (DOE) Joint Genome Institute (JGI)³⁸ and deposited in the European Molecular Biology Laboratory (EMBL)/GenBank/ DNA Data Bank of Japan (DDBJ) database³⁸. The genome size is 7.1 Mbp, with a total of ~5754 predicted protein-coding open-reading frames³⁸. Availability of the full genome sequence should lead to a rapid increase in our knowledge of functional genes involved in growth, development, and metabolic processes in this organism. Thus, key gene networks supporting the different metabolic processes in various cell types are now within reach for identification.

Two enzyme systems of hydrogenase and nitrogenase play a major role in algal hydrogen production^{106, 111}. *T. variabilis* hydrogenase and nitrogenase enzymes and genes encoding these proteins have been the subject of much research¹¹². Hydrogenase enzymes play the main role in photolysis-dependent hydrogen production as well as hydrogen utilisation. There are two kinds of hydrogenase enzymes: (i) S-Fe hydrogenase (known as reversible or soluble hydrogenase), encoded by *hoxEFUYH* genes¹¹³⁻¹¹⁵, function in generating molecular hydrogen to reduce NAD required for CO₂ absorption in the ribulose biphosphate cycle and (ii) Ni- Fe hydrogenase (known as uptake or membrane hydrogenase), encoded by *hupSL* genes¹¹⁶, which catalyses electron transfer from molecular hydrogen to the respiratory chain. Theoretically, efficiency of H₂ production in direct photolysis is high (40.1%) but it is not practically feasible due to the high inhibitory effects of O₂ on the reversible hydrogenase^{103, 106, 117}.

Since nitrogenase enzymes are sensitive to oxygen, this requires nitrogenase-catalysed H₂-generating reactions to be separated from oxygen-containing environments. Therefore, indirect photolysis evolved, in which hydrogen is generated in two separate stages. In the first stage, electrons derived from water splitting are consumed in CO₂ fixation to sugars. Thus, CO₂ is stored in vegetative cells as

carbohydrate and then transported to heterocysts, where the sugar is used as an electron donor^{118, 119}. In the second stage, breakdown of the sugar releases electrons used to produce hydrogen by nitrogenase enzymes¹⁰³. In the heterocysts of *T. variabilis*, groups of *nif* and *vnf* genes are expressed to give rise to metal ion-dependent nitrogenase, which is responsible for nitrogen fixing and generates hydrogen as a by-product of the process. There are three main gene clusters encoding nitrogenases, depending on distinct prosthetic groups: Mo-Fe (*nif1* and *nif2* genes), V-Fe (*vnf* genes) and Fe-Fe (*anf* genes) nitrogenases. Mo-Fe nitrogenases are encoded by (i) groups of *nif1* genes, including *nifBSUHDKENXW*^{20, 120-123} and three open reading frames expressed under diazotrophic conditions, and (ii) *nif2* genes expressed under anoxic conditions. The latter are essentially the same (*nifBSUHDKENXW*) genes with the exception of a few differences, such as the fusion of *nifX* and *nifEN* genes into a single open reading frame¹²⁴. V-Fe nitrogenase contains $\alpha\beta\delta$ -subunits and scaffold proteins encoded by *vnfDKGENH*¹²⁵ which is expressed under diazotrophic conditions in the absence of Mo, with or without V. The third nitrogenase known as Fe-Fe nitrogenase is encoded by *anfHDKG*¹²⁶. Further reviews and reports of *T. variabilis* nitrogenase gene expression and heterocyst metabolism have been published^{124, 125, 127-131}. Although indirect photolysis is more feasible than direct photolysis, the H₂ production efficiency is lower (16.3%) due to the multiphase process and ATP consumption required to drive the nitrogenase activity¹⁰³.

Other options based on fermentation of biomass have been developed. Hey *et al*¹³² reviewed key factors affecting photo and dark fermentation, including substrates, inoculum density, and environmental conditions. Light fermentation is similar to indirect photolysis. During indirect photolysis, endogenous organic compounds are used to produce hydrogen, while in photofermentation exogenous organic compounds are consumed for use in generating electrons. Thus, fermentation is considered the most promising method for microbial H₂ production¹⁰³. Unlike light driven processes, during dark fermentation organic substrates are used both as energy and electron sources. Bartacek *et al*¹³³ summarized fundamentals of fermentative H₂ generation in different microorganisms. Although anaerobic bacteria are mostly used in dark fermentation, it was reported that the combination of photo and dark fermentation of photosynthetic bacteria and algae could significantly enhance the bio-hydrogen productivity¹³⁴. To the best of our knowledge, there are no reports on application of *T. variabilis* in such a hybrid system. Thus, further research on metabolic pathways of dark-driven hydrogen generation using *T. variabilis* is required.

6.4 Factors affecting H₂ generation in *T. variabilis*

Maximal H₂ production from algal cultures can be achieved through optimisation of key physicochemical factors with a regulatory influence on the generation capacity. High light intensity inhibits H₂ production rate due to up-regulation of photosynthesis, which enhances oxygen accumulation and consequent inhibition of nitrogenases^{104, 135}. Darkness, on the other hand, can block H₂ production via depletion of carbohydrate reserves, which triggers a deficiency of energy and electrons needed for nitrogenase activity^{118, 136}. The negative effects of excessive oxygen can be overcome through degassing or generating a partial vacuum¹³⁷, sparging with argon gas¹³⁸ and temporal separation of the photosynthetic phase from the H₂ generation stage⁴⁸. Based on the 2-stage system for H₂ production, a flat panel photo-bioreactor was found to increase biomass production in the first stage with N-replete medium, and the H₂ generating in

the second stage with N-depleted medium sparging with pure Ar⁴⁸.

The presence of a nitrogen source in the growth medium severely decreases H₂ production, particularly when the nitrogenase enzyme machinery is the dominant H₂-producing complex. This is because inorganic nitrogen suppresses differentiation of heterocysts, thereby removing heterocyst-dependent generation. Although H₂ could be generated as a by-product of N-fixation utilizing atmospheric nitrogen as noted earlier (section 5.3) and the frequency of heterocysts increases in N-free growth medium¹³⁹, the innate biochemical inefficiencies of this process reduce H₂ gas production. About 75% of electrons are diverted away from H₂ synthesis to ammonium reduction, and this loss can be averted by the absence of atmospheric nitrogen¹¹⁸.

H₂ production rate diminishes with decreasing pH¹⁴⁰. A study focusing on pH effects demonstrated that algal growth rate was 3 times higher in the presence of Mo- or V-nitrogenase than in the presence of Fe- nitrogenase and maximum H₂ production was attained in the presence of V-nitrogenase in a wide range of alkaline cultures (pH 7-9)¹⁴¹. Mo and V are required as essential components in Mo-nitrogenase and V-nitrogenase based H₂ production, respectively, provided in BG and AA growth media^{118, 142}. Although Mo significantly increases H₂ production rate, it has a specific upper limit (1.6 mM) beyond which further increases do not influence productivity¹¹⁸. The maximum light to hydrogen energy conversion efficiencies were achieved in Allen-Arnon media, where a larger heterocyst frequency was noted⁴⁸. Temperature is another parameter affecting H₂ photo-production. A short-term thermal stress (30-36 °C) enhances nitrogenase activity of *T. variabilis*¹⁴³. The results of studying CO₂ concentration, as a carbon source, demonstrated that an optimum initial CO₂ molar fraction of 0.05 is required for maximum biomass production⁵⁶. Additionally, inclusion of a carbon source, such as glucose, gives rise to further hydrogen production¹¹⁸.

6.5 Genetic approach for increasing *T. variabilis* H₂ production

Manipulation of the parameters dealt with in the preceding section (e.g., pH, temperature, and nutrients) can be used to increase hydrogen production by algal systems. However, substantial research efforts have been directed at other key targets to enhance hydrogen yield. For example, the reversible hydrogenase enzyme system, which can consume as well as generate hydrogen, is an obvious target for inactivation using genetic approaches. Similarly, the uptake hydrogenase, which only consumes H₂ can be targeted for gene deletion. Accordingly, a *T. variabilis* loss-of-function mutant (AVM13) in the *hupSL* gene, which encodes the uptake hydrogenase, had at least a 3-fold increase in H₂ production when compared to the wildtype¹⁴⁴. Nitroso-guanidine mutagenesis was used to generate PK84 and PK17R *T. variabilis* mutants with loss-of-function mutations impairing uptake hydrogenase in both mutants and an additional impairment of bidirectional (reversible) hydrogenase in PK84¹⁴⁵. When compared to the wildtype, H₂ production in PK17R and PK84 was increased 1.4-fold and 4.3-fold, respectively¹⁴⁶. Notably, while wild-type strains produce appreciable amounts of H₂ only in an argon atmosphere (to preclude O₂), the mutant PK84 produced H₂ in a CO₂ atmosphere, which is more desirable from a process scale-up perspective¹⁴⁷. While mutants are showing great promise (Table 2), environmental concerns about the release of genetically modified organisms exist¹⁴⁸. Since open biomass production systems are likely to be used in combined phycoremediation and bioenergy generation, use of mutant algae is inevitably problematic.

6.6 Feasibility of integrated biofuel production and wastewater treatment

In order to make the application of microalgae system feasible at industrial scale, it is vital to use sunlight as an energy source to significantly reduce the operation costs, relative to using artificial lighting. The mutant form of *T. variabilis* has demonstrated viable growth in ambient outdoor conditions¹⁴⁹⁻¹⁵¹. It was shown that in such conditions, this species can prolong hydrogen generation without medium refreshment for 40 days¹⁵¹. Also the feasibility of growing the wildtype of this species in wastewater had promising results⁹. Although there are some pilot studies concentrated on bioenergy production and/or wastewater treatment using *T. variabilis*, there still is lack of specific research examining performance at larger industrial scale. However, several research groups have demonstrated that under certain circumstances, large-scale H₂-photo production using microalgae is viable¹⁵². Previously, scaling up the algal systems for bioenergy production was investigated from different techno-economic and socio-environmental angles considering various parameters, such as water and energy consumption, cultivation and harvesting technologies, and associated costs and savings⁸⁴. Open raceway ponds with lower capital and operational costs, in comparison to PBRs, represent an attractive option for commercial large-scale bioenergy production¹⁵³. It was demonstrated that this technology provides a "win-win strategy"¹⁵⁴, having remarkable productivity with an average capacity of 40-70 tons biomass.ha⁻¹.year⁻¹ during bioremediation process, which can be coupled with biorefinery¹⁵³. Results of a feasibility study for integration of wastewater treatment and biogas production using microalgae systems indicated positive benefits on payback period and internal rate of return (*IRR*) for biofuel production when considering environmental revenue and cost-savings, such as bio-products income, carbon credits and wastewater treatment¹⁵⁵. The same strategy was reported as a feasible solution for using algae and bacteria cocultures to couple bioremediation and biofuel production¹⁵⁶. Economic feasibility of commercial raceway pond use in wastewater treatment and biogas production should consider the pond size, electricity requirements, and thermal capacity for bio-methane production¹⁵⁷. In such a system, the required CO₂ and thermal energy could be met by generation via combined heat and power production (305 kg CO₂.d⁻¹ and 488 kWh.d⁻¹) coupled with biogas production system¹⁵⁷. Therefore, the limited information on feasibility of bioenergy production using microalgal systems points to a beneficial outcome with significantly reduced energy consumption and costs if the system integrates with bioremediation to cover nutrient (40–100 kg of N and 3-12 kg of P) and water (11–13 ML.ha⁻¹.year⁻¹) requirements¹⁵³.

7 Challenges and opportunities

There are several studies on various aspects of *T. variabilis* biology, cultivation, and application to phycoremediation. Here, we have provided a coherent synthesis of the literature by presenting how surveying the interrelation between different studies could develop a route to industrial application of *T. variabilis* in bioremediation and production of sustainable bioenergy, discussing challenges, opportunities and research gaps as bellow:

- 1) Bioremediation performance of *T. variabilis* illustrated promising results in bioabsorption of nutrients and heavy metals such as Cu, Zn, and Ni. Remediation of different metals may influence gene expression and enzymatic function. This would be more important when the biomass harvested from remediation process

is used directly for bioenergy production. Thus, research into the effects of remediation on metabolic pathways, which control bioenergy generation, in this species needs to be undertaken.

- 2) There are some pilot studies showing the utility of *T. variabilis* in bioremediation using artificially constituted “wastewaters”. However, actual wastewaters may contain unknown hazardous compounds and allelochemicals, which can negatively influence algal growth. For example, nTiO₂, Microcystins, Harmane and FischerellinB restrict application of algae in wastewater streams containing such pollutants. Therefore, it is necessary to examine actual biomass generation in different real wastewater streams for scaling up phycoremediation at industrial scale for biorefining purposes.
- 3) The application of *T. variabilis* for biodiesel and bio-ethanol production has not been widely studied. Although *T. variabilis* is not considered an oleaginous species, environmental stress can induce cells to produce significant amounts of lipid. The molecular genetic basis for the enhanced lipid synthesis requires further research. Also, bio-ethanol production using this organism is not viable due to the challenge of polysaccharide loss together with associated costly and high energy-consuming technologies required for the process. Therefore, more efficient solutions are required.
- 4) Biogas generation using this alga is promising, particularly when coupled to wastewater treatment. In order to improve bio-methane yield, first it is necessary to evaluate the effect of the bioremediation process on algal cell composition, protein profile and biomass bio-degradability. This would provide valuable information to boost the initial hydrolysis processes via applying the most appropriate pretreatment technology. Also, applying co-digestion technologies can enhance anaerobic digestion kinetics and, from an economic aspect, there are still several improvement methods such as optimizing the proportions of microalgal cells for declining the anaerobic digestion hydraulic retention times which need further investigation.
- 5) Bio-hydrogen production using *T. variabilis* is focused on light-driven methods and the potential of using this species in dark-driven hybrid systems, particularly in combination with dark fermentation, has not yet been surveyed. Although the challenge of oxygen inhibitory effects on hydrogen generation can be averted by degassing, partial vacuum, sparging Ar, applying two-staged H₂ generation method or utilising mutants of *T. variabilis*, presented solutions are costly and carry environmental risks. Therefore, still further research is necessary to develop cost-effective and environmentally friendly alternatives for robust bio-hydrogen production systems.

8 Future prospects

In this review, we have sought to collate the wide literature on *Trichormus variabilis*, bringing data together from many different aspects, including morphology focusing on different types of cell differentiation and physiology, together with a review of growth parameters and conditions where the species may be used in remediation of wastewaters coupled with production of different forms of liquid and gaseous bioenergy, such as biodiesel, bio-ethanol, bio-methane and bio-hydrogen. We attempted to distil the key information available on *T. variabilis* on phycoremediation and bioenergy production to enable further research on optimization of robust technologies for supplying clean water

and sustainable energy resources. Trends in research of *T. variabilis* during the last decades has focused on the main aspects of (i) biology, (ii) nutrient removal and (iii) bioenergy production, and considered the effects of several stress factors on growth and lipid production. There is a promising path for future application of this species even though there are still major environmental and economic issues which need to be resolved to optimise *T. variabilis* for industrial usage. The valuable potential of this microorganism arising from the differentiation of vegetative cells into other cell types for adaptive responses to different environmental conditions provides a great opportunity to develop platforms for commercial exploitation of this species. It is important to elucidate the mechanisms of cell differentiation and metabolic changes associated with exposure to stress.

Acknowledgements

The authors are grateful to Christine Richardson (Department of Biosciences, Durham University) for assistance with transmission electron microscopy.

Competing interests

The authors have no competing interests.

References

1. Conti J, Holtberg P, Diefenderfer J, LaRose A, Turnure JT, Westfall L. World energy demand and economic outlook, Chapter 1. *International Energy Outlook 2016 With Projections to 2040*. Energy Information Administration (EIA), Office of Energy Analysis., Washington. pp. 7-17 (2016).
2. Oswald W. The high-rate pond in waste disposal. *Developments in Industrial Microbiology*. **4**:112-25(1963).
3. Olguin xn, Eugenia J. Phycoremediation: key issues for cost-effective nutrient removal processes. *Biotechnology Advances*. **22**:81-91(2003).
4. Esquivel-Hernández DA, Ibarra-Garza IP, Rodríguez-Rodríguez J, Cuéllar-Bermúdez SP, Rostro-Alanis MdJ, Alemán-Nava GS, et al. Green extraction technologies for high-value metabolites from algae: a review. *Biofuels, Bioproducts and Biorefining*. **11**:215-31(2017).
5. Chekroun KB, Baghour M. The role of algae in phytoremediation of heavy metals: a review. *Journal of Materials and Environmental Science*. **4**:873-80(2013).
6. Brennan L, Owende P. Biofuels from microalgae-A review of technologies for production, processing, and extractions of biofuels and co-products. *Renewable and Sustainable Energy Reviews*. **14**:557-77(2010).
7. Greenwell HC, Laurens LML, Shields RJ, Lovitt RW, Flynn KJ. Placing microalgae on the biofuels priority list: a review of the technological challenges. *Journal of The Royal Society Interface*. **7**:703-26(2010).
8. Wijffels RH, Barbosa MJ, Eppink MH. Microalgae for the production of bulk chemicals and biofuels. *Biofuels, Bioproducts and Biorefining: Innovation for a sustainable economy*. **4**:287-95(2010).
9. El-Bestawy E. Treatment of mixed domestic-industrial wastewater using cyanobacteria. *Journal of Industrial Microbiology and Biotechnology*. **35**:1503-16(2008).

- 1 10.Noel SD, Rajan M. Cyanobacteria as a potential source of phycoremediation from
2 textile industry effluent. *Journal of Microbiology and Biotechnology Research*. **4**:35-
3 3(2014).
- 4 11.Talukder A, Mahmud S, Lira S, Aziz M. Phycoremediation of Textile Industry
5 Effluent by Cyanobacteria (*Nostoc muscorum* and *Anabaena variabilis*). *Bioresearch*
6 *Communications*. **1**:124-7(2015).
- 7 12.Khetkorn W, Rastogi RP, Incharoensakdi A, Lindblad P, Madamwar D, Pandey A, et
8 al. Microalgal hydrogen production—A review. *Bioresource technology*. **243**:1194-
9 206(2017).
- 10 13.Markov SA, Protasov ES, Bybin VA, Eivazova ER, Stom DI. Using immobilized
11 cyanobacteria and culture medium contaminated with ammonium for H₂ production in a
12 hollow-fiber photobioreactor. *International Journal of Hydrogen Energy*. **40**:4752-
13 7(2015).
- 14 14.Ismail G, Abo-Hamad SAE. Effect of different *Anabaena variabilis* (Kütz)
15 treatments on some growth parameters and physiological aspects of *Hordeum vulgare*
16 *L.* and *Trigonella foenum-graecum L.* *Egyptian Journal of Botany*. **57**:507-16(2017).
- 17 15.Guiry MD, Guiry GM. AlgaeBase. *World-wide electronic publication*, National
18 University of Ireland, Galway. <http://www.algaebase.org>(2019).
- 19 16.Horton T, Kroh A, Ah Yong S, Bailly N, Boyko CB, Brandão SN, et al. World
20 Register of Marine Species (WoRMS). <http://www.marinespecies.org>(2019).
- 21 17.Inkscape Team. *Inkscape*: A vector drawing tool. <http://www.inkscape.org>(2004).
- 22 18.Abedi S, Astaraei FR, Ghobadian B, Tavakoli O, Jalili H, Greenwell HC, et al.
23 Decoupling a novel *Trichormus variabilis*-*Synechocystis sp.* interaction to boost
24 phycoremediation. *Scientific reports*. **9**:2511(2019).
- 25 19.Rajaniemi P, Hrouzek P, Kastovska K, Willame R, Rantala A, Hoffmann L, et al.
26 Phylogenetic and morphological evaluation of the genera *Anabaena*, *Aphanizomenon*,
27 *Trichormus* and *Nostoc* (Nostocales, Cyanobacteria). *International Journal of*
28 *Systematic and Evolutionary Microbiology* **55**:11-26(2005).
- 29 20.Gladkikh AS, Belykh OI, Klimenkov IV, Tikhonova IV. Nitrogen-fixing
30 cyanobacterium *Trichormus variabilis* of the lake Baikal phytoplankton.
31 *Mikrobiologiya*. **77**:814-22(2008).
- 32 21.Kützing FT. *Phycologia Generalis, Clastotype for Anabaena variabilis* Kütz.
33 (*Nostocaceae*). Smithsonian Institution, National Museum of Natural History,
34 Department of Botany, : Encyclopedia of life, US 139159; (1839).
- 35 22.Gkelis S, Ourailidis I, Panou M, Pappas N. Cyanobacteria of Greece: an annotated
36 checklist. *Biodiversity Data Journal*. **4**:e10084(2016).
- 37 23.Shariatmadari Z, Riahi H. A taxonomic study on soil taxa of apoheterocytic
38 cyanoprokaryota from Nostocaceae family in Iran. *The Iranian Journal of Botany*.
39 **19**:143-52(2013).
- 40 24.Prasanna R, Kumar R, Sood A, Prasanna BM, Singh PK. Morphological,
41 physiochemical and molecular characterization of *Anabaena* strains. *Microbiological*
42 *Research*. **161**:187-202(2006).
- 43 25.Komárek J, Johansen JR. Chapter 4 - Filamentous Cyanobacteria. In: Sheath RG,
44 Kociolek JP, editors. *Freshwater Algae of North America (Second Edition)*. Academic
45 Press., Boston, pp. 135-235 (2015).
- 46 26.Komarek JaK, Anagnostidis,. Modern approach to the classification system of
47 Cyanophytes 4-Nostocales. *Arch Hydrobiol Suppl* 82(3), *Algological Studies/Archiv für*
48 *Hydrobiologie, Supplement Volumes*. **56**:247-345(1989).

27. Andreja K, Kozlikova-Zapomelova E, Mareš J, Řeháková K. A detailed morphological, phylogenetic and ecophysiological analysis of four benthic *Anabaena* (Nostocales, Cyanobacteria) strains confirms deep heterogeneity within the genus. *Fottea, Olomouc*. **15**:191-202(2015).
28. R Y Stanier a, Bazine GC. Phototrophic Prokaryotes: The Cyanobacteria. *Annual Review of Microbiology*. **31**:225-74(1977).
29. Wolgemuth CW, Oster G. The junctional pore complex and the propulsion of bacterial cells. *Journal of molecular microbiology and biotechnology*. **7**:72-7(2004).
30. Adams DG, Duggan PS. Tansley Review No. 107. Heterocyst and akinete differentiation in cyanobacteria. *The New Phytologist*. **144**:3-33(1999).
31. Park JJ, Sigal LY, Wolk CP, Vieille C. Cell-specific gene expression in *Anabaena variabilis* grown phototrophically, mixotrophically, and heterotrophically. *BMC Genomics*. **14**:759(2013).
32. Lynn ME, Bantle JA, Ownby JD. Estimation of gene expression in heterocysts of *Anabaena variabilis* by using DNA-RNA hybridization. *Journal of bacteriology*. **167**:940-6(1986).
33. Flores E, Pernil R, Muro-Pastor AM, Mariscal V, Maldener I, Lechno-Yossef S, et al. Septum-localized protein required for filament integrity and diazotrophy in the heterocyst-forming cyanobacterium *Anabaena* sp. strain PCC 7120. *Journal of bacteriology*. **189**:3884-90(2007).
34. Sadava DE, Orians GH, Heller HC, Hillis DM, Purves WK. The Evolution of Diversity: Bacteria, Archaea and Viruses. *Life: The Science of Biology*. Palgrave Macmillan., Basingstok, pp. 532-3 (2013).
35. Maldener I, Summers ML, Sukenik A. Cellular differentiation in filamentous cyanobacteria. *The Cell Biology of Cyanobacteria*. **15**:263-91(2014).
36. Perez R, Wörmer L, Sass P, Maldener I. A highly asynchronous developmental program triggered during germination of dormant akinetes of filamentous diazotrophic cyanobacteria. *FEMS Microbiology Ecology*. **94**:fix131(2018).
37. Zhu L. Microalgal culture strategies for biofuel production: a review. *Biofuels, Bioproducts and Biorefining*. **9**:801-14(2015).
38. Thiel T, Pratte BS, Zhong J, Goodwin L, Copeland A, Lucas S, et al. Complete genome sequence of *Anabaena variabilis* ATCC 29413. *Standards in Genomic Sciences*. **9**:562-73(2014).
39. Mannan RM, Pakrasi HB. Dark heterotrophic growth conditions result in an increase in the content of photosystem II units in the filamentous cyanobacterium *Anabaena variabilis* ATCC 29413. *Plant Physiology*. **103**:971-7(1993).
40. Collins CD, Boylen CW. Physiological responses of *Anabaena variabilis* (Cyanophyceae) to instantaneous exposure to various combinations of light intensity and temperature. *Journal of Phycology*. **18**:206-11(1982).
41. Yoon JH, Sim SJ, Kim MS, Park TH. High cell density culture of *Anabaena variabilis* using repeated injections of carbon dioxide for the production of hydrogen. *International Journal of Hydrogen Energy*. **27**:1265-70(2002).
42. Stanier RY, Kunisawa R, Mandel M, Cohen-Bazire G. Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriological Reviews*. **35**:171-205(1971).
43. Allen MB, Arnon DI. Studies on nitrogen-fixing blue-green algae. I. growth and nitrogen fixation by *Anabaena cylindrica* Lemm. *Plant Physiology*. **30**:366-72(1955).

1 44.Starr RC, Zeikus JA. UTEX—the culture collection of algae at the University of
2 Texas at Austin 1993 list of cultures. *Journal of phycology*. **29**:1-106(1993).

3 45.Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. Generic
4 assignments, strain histories and properties of pure cultures of cyanobacteria.
5 *Microbiology*. **111**:1-61(1979).

6 46.Carmichael WW, Gorham PR. An improved method for obtaining axenic clones of
7 planktonic blue-green algae. *Journal of Phycology*. **10**:238-40(1974).

8 47.Fogg G. Growth and heterocyst production in *Anabaena cylindrica* lemm.: II. in
9 relation to carbon and nitrogen metabolism. *Annals of botany*. **13**:241-59(1949).

10 48.Berberoglu H, Jay J, Pilon L. Effect of nutrient media on photobiological hydrogen
11 production by *Anabaena variabilis* ATCC 29413. *International Journal of Hydrogen*
12 *Energy*. **33**:1172-84(2008).

13 49.Kaplan A, Badger MR, Berry JA. Photosynthesis and the intracellular inorganic
14 carbon pool in the bluegreen alga *Anabaena variabilis*: response to external CO₂
15 concentration. *Planta*. **149**:219-26(1980).

16 50.Troshina OY, Serebryakova LT, Lindblad P. Induction of H₂-uptake and nitrogenase
17 activities in the cyanobacterium *Anabaena variabilis* ATCC 29413: effects of hydrogen
18 and organic substrate. *Current Microbiology*. **33**:11-5(1996).

19 51.Salleh SF, Kamaruddin A, Uzir MH, Mohamed AR, Shamsuddin AH. Modeling the
20 light attenuation phenomenon during photoautotrophic growth of *A. variabilis* ATCC
21 29413 in a batch photobioreactor. *Journal of Chemical Technology and Biotechnology*.
22 **92**:358-66(2017).

23 52.Fallowfield H, Osborne B. Growth and light absorptance of Cyanobacteria and
24 Chlorophyceae with particular reference to *Anabaena variabilis* and *Scenedesmus*
25 *obliquus*. *British Phycological Journal*. **20**:27-41(1985).

26 53.Haury JF, Spiller H. Fructose uptake and influence on growth of and nitrogen
27 fixation by *Anabaena variabilis*. *Journal of bacteriology*. **147**:227-35(1981).

28 54.Mimuro M, Fujita Y. Estimation of chlorophyll a distribution in the photosynthetic
29 pigment systems I and II of the blue-green alga *Anabaena variabilis*. *Biochimica et*
30 *Biophysica Acta (BBA)-Bioenergetics*. **459**:376-89(1977).

31 55.Nozue S, Katayama M, Terazima M, Kumazaki S. Comparative study of thylakoid
32 membranes in terminal heterocysts and vegetative cells from two cyanobacteria,
33 *Rivularia M-261* and *Anabaena variabilis*, by fluorescence and absorption spectral
34 microscopy. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*. **1858**:742-9(2017).

35 56.Berberoglu H, Barra N, Pilon L, Jay J. Growth, CO₂ consumption and H₂ production
36 of *Anabaena variabilis* ATCC 29413-U under different irradiances and CO₂
37 concentrations. *J Appl Microbiol*. **104**:105-21(2008).

38 57.Samorì G, Samorì C, Guerrini F, Pistocchi R. Growth and nitrogen removal capacity
39 of *Desmodesmus communis* and of a natural microalgae consortium in a batch culture
40 system in view of urban wastewater treatment: Part I. *Water Research*. **47**:791-
41 801(2013).

42 58.El-Sheekh MM, El-Shouny WA, Osman MEH, El-Gammal EWE. Growth and heavy
43 metals removal efficiency of *Nostoc muscorum* and *Anabaena subcylindrica* in sewage
44 and industrial wastewater effluents. *Environmental Toxicology and Pharmacology*.
45 **19**:357-65(2005).

46 59.González LE, Cañizares RO, Baena S. Efficiency of ammonia and phosphorus
47 removal from a colombian agroindustrial wastewater by the microalgae *Chlorella*
48 *vulgaris* and *Scenedesmus dimorphus*. *Bioresource Technology*. **60**:259-62(1997).

1 60.Kong Q-x, Li L, Martinez B, Chen P, Ruan R. Culture of microalgae
2 *Chlamydomonas reinhardtii* in wastewater for biomass feedstock production. *Applied*
3 *Biochemistry and Biotechnology*. **160**:9(2009).

4 61.Wang L, Min M, Li Y, Chen P, Chen Y, Liu Y, et al. Cultivation of green algae
5 *Chlorella sp.* in different wastewaters from municipal wastewater treatment plant.
6 *Applied Biochemistry and Biotechnology*. **162**:1174-86(2010).

7 62.Caporgno MP, Taleb A, Olkiewicz M, Font J, Pruvost J, Legrand J, et al. Microalgae
8 cultivation in urban wastewater: Nutrient removal and biomass production for biodiesel
9 and methane. *Algal Research*. **10**:232-9(2015).

10 63.Worku A, Sahu O. Reduction of heavy metal and hardness from ground water by
11 algae. *Journal of Applied & Environmental Microbiology*. **2**:86-9(2014).

12 64.Zhou JL, Huang PL, Lin RG. Sorption and desorption of Cu and Cd by macroalgae
13 and microalgae. *Environmental Pollution*. **101**:67-75(1998).

14 65.Chevalier P, Proulx D, Lessard P, Vincent WF, de la Noüe J. Nitrogen and
15 phosphorus removal by high latitude mat-forming cyanobacteria for potential use in
16 tertiary wastewater treatment. *Journal of Applied Phycology*. **12**:105-12(2000).

17 66.Kaewsarn P. Biosorption of copper(II) from aqueous solutions by pre-treated
18 biomass of marine algae *Padina sp.* *Chemosphere*. **47**:1081-5(2002).

19 67.Freitas OMM, Martins RJE, Delerue-Matos CM, Boaventura RAR. Removal of
20 Cd(II), Zn(II) and Pb(II) from aqueous solutions by brown marine macro algae: Kinetic
21 modelling. *Journal of Hazardous Materials*. **153**:493-501(2008).

22 68.Soeprobawati TR, Hariyati R. Bioaccumulation of Pb, Cd, Cu, and Cr by
23 *Porphyridium cruentum* (SF Gray) Nägeli. *International Journal of Marine Science*.
24 **3**:(2013).

25 69.Jia H, Yuan Q. Removal of nitrogen from wastewater using microalgae and
26 microalgae–bacteria consortia. *Cogent Environmental Science*. **2**:1275089(2016).

27 70.Parameswari E, Lakshmanan A, Thilagavathi T. Phycoremediation of heavy metals
28 in polluted water bodies. *Electronic Journal of Environmental, Agricultural and Food*
29 *Chemistry*. **9**:808-14(2010).

30 71.Hirooka T, Akiyama Y, Tsuji N, Nakamura T, Nagase H, Hirata K, et al. Removal of
31 hazardous phenols by microalgae under photoautotrophic conditions. *Journal of*
32 *Bioscience and Bioengineering*. **95**:200-3(2003).

33 72.Bauer N, Blaschke U, Beutler E, Gross E, Jenett-Siems K, Siems K, et al. Seasonal
34 and interannual dynamics of polyphenols in *Myriophyllum verticillatum* and their
35 allelopathic activity on *Anabaena variabilis*. *Aquatic Botany*. **91**:110-6(2009).

36 73.Cherchi C, Miljkovic M, Diem M, Gu AZ. nTiO₂ induced changes in intracellular
37 composition and nutrient stoichiometry in primary producer-cyanobacteria. *The Science*
38 *of the total environment*. **512**:345-52(2015).

39 74.Cherchi C, Chernenko T, Diem M, Gu AZ. Impact of nano titanium dioxide
40 exposure on cellular structure of *Anabaena variabilis* and evidence of internalization.
41 *Environmental Toxicology and Chemistry*. **30**:861-9(2011).

42 75.Cherchi C, Gu AZ. Impact of Titanium Dioxide Nanomaterials on Nitrogen Fixation
43 Rate and Intracellular Nitrogen Storage in *Anabaena variabilis*. *Environmental Science*
44 *& Technology*. **44**:8302-7(2010).

45 76.Kodani S, Imoto A, Mitsutani A, Murakami M. Isolation and identification of the
46 antialgal compound, harmane (1-methyl- β -carboline), produced by the algicidal
47 bacterium, *Pseudomonas sp. K44-1*. *Journal of applied phycology*. **14**:109-14(2002).

- 1 77.Gross E. Allelopathy in Benthic and Littoral Areas: Case Studies on Allelochemicals
2 from Benthic Cyanobacteria and Submersed Macrophytes. In: Inderjit, Dakshini KMM,
3 Foy CL, editors. *Principles and Practices in Plant Ecology*. CRC Press., Boca Raton,
4 Florida, pp. 179-99 (1999).
- 5 78.Chia MA, Jankowiak JG, Kramer BJ, Goleski JA, Huang I-S, Zimba PV, et al.
6 Succession and toxicity of *Microcystis* and *Anabaena* (Dolichospermum) blooms are
7 controlled by nutrient-dependent allelopathic interactions. *Harmful algae*. **74**:67-
8 77(2018).
- 9 79.Bártová K, Hilscherová K, Babica P, Marsálek B. Extract of *Microcystis* water
10 bloom affects cellular differentiation in filamentous cyanobacterium *Trichormus*
11 *variabilis* (Nostocales, Cyanobacteria). *Journal of applied phycology*. **23**:967-73(2011).
- 12 80.Singh NK, Upadhyay AK, Rai UN. Algal technologies for wastewater treatment and
13 biofuels production: an integrated approach for environmental management. In: Gupta
14 SK, Malik A, Bux F, editors. *Algal Biofuels: Recent Advances and Future Prospects*.
15 Springer International Publishing., Cham, pp. 97-107 (2017).
- 16 81.Rawat I, Ranjith Kumar R, Mutanda T, Bux F. Dual role of microalgae:
17 Phycoremediation of domestic wastewater and biomass production for sustainable
18 biofuels production. *Applied Energy*. **88**:3411-24(2011).
- 19 82.Cho S, Luong TT, Lee D, Oh Y-K, Lee T. Reuse of effluent water from a municipal
20 wastewater treatment plant in microalgae cultivation for biofuel production.
21 *Bioresource Technology*. **102**:8639-45(2011).
- 22 83.Kligerman DC, Bouwer EJ. Prospects for biodiesel production from algae-based
23 wastewater treatment in Brazil: A review. *Renewable and Sustainable Energy Reviews*.
24 **52**:1834-46(2015).
- 25 84.Xin C, Addy MM, Zhao J, Cheng Y, Cheng S, Mu D, et al. Comprehensive techno-
26 economic analysis of wastewater-based algal biofuel production: A case study.
27 *Bioresource technology*. **211**:584-93(2016).
- 28 85.Chen Y, Xu C, Vaidyanathan S. Microalgae: A robust “green bio-bridge” between
29 energy and environment. *Critical reviews in biotechnology*. **38**:351-68(2018).
- 30 86.Atabani A, Silitonga A, Ong H, Mahlia T, Masjuki H, Badruddin IA, et al. Non-
31 edible vegetable oils: a critical evaluation of oil extraction, fatty acid compositions,
32 biodiesel production, characteristics, engine performance and emissions production.
33 *Renewable and sustainable energy reviews*. **18**:211-45(2013).
- 34 87.Hajinezhad A, Abedi S, Ghobadian B, Noorollahi Y. Biodiesel production from
35 Norouzak (*Salvia lerifolia*) seeds as an indigenous source of bio fuel in Iran using
36 ultrasound. *Energy Conversion and Management*. **99**:132-40(2015).
- 37 88.Vargas M, Rodriguez H, Moreno J, Olivares H, Del Campo J, Rivas J, et al.
38 Biochemical composition and fatty acid content of filamentous nitrogen-fixing
39 cyanobacteria. *Journal of phycology*. **34**:812-7(1998).
- 40 89.Han F, Pei H, Hu W, Jiang L, Cheng J, Zhang L. Beneficial changes in biomass and
41 lipid of microalgae *Anabaena variabilis* facing the ultrasonic stress environment.
42 *Bioresource Technology*. **209**:16-22(2016).
- 43 90.Rodolfi L, Chini Zittelli G, Bassi N, Padovani G, Biondi N, Bonini G, et al.
44 Microalgae for oil: Strain selection, induction of lipid synthesis and outdoor mass
45 cultivation in a low-cost photobioreactor. *Biotechnology and bioengineering*. **102**:100-
46 12(2009).

- 1 91.Han F, Pei H, Hu W, Song M, Ma G, Pei R. Optimization and lipid production
2 enhancement of microalgae culture by efficiently changing the conditions along with
3 the growth-state. *Energy Conversion and Management*. **90**:315-22(2015).
- 4 92.Wang ZT, Ullrich N, Joo S, Waffenschmidt S, Goodenough U. Algal lipid bodies:
5 stress induction, purification, and biochemical characterization in wild-type and
6 starchless *Chlamydomonas reinhardtii*. *Eukaryotic Cell*. **8**:1856-68(2009).
- 7 93.Pyo DJ,Method for producing ethanol using freshwater blue-green algae .Korea
8 patent WO2010095895A9 (2010).
- 9 94.Pyo D, Kim T, Yoo J. Efficient extraction of bioethanol from freshwater
10 cyanobacteria using supercritical fluid pretreatment. *Bull Korean Chem Soc*. **34**:379-
11 83(2013).
- 12 95.Weiland P. Biogas production: current state and perspectives. *Applied microbiology*
13 *and biotechnology*. **85**:849-60(2010).
- 14 96.Mussnug JH, Klassen V, Schlüter A, Kruse O. Microalgae as substrates for
15 fermentative biogas production in a combined biorefinery concept. *Journal of*
16 *biotechnology*. **150**:51-6(2010).
- 17 97.Petrova E, Egorova M, Piskunkova N, Kozhevin P, Netrusov A, Tsavkelova E.
18 Anaerobic cellulolytic microbial communities decomposing the biomass of *Anabaena*
19 *variabilis*. *Microbiology*. **86**:745-52(2017).
- 20 98.Gassanova L, Netrusov AI, Teplyakov V, Modigell M. Fuel gases from organic
21 wastes using membrane bioreactors. *Desalination*. **198**:56-66(2006).
- 22 99.Magdalena J, Ballesteros M, González-Fernandez C. Efficient anaerobic digestion of
23 microalgae biomass: proteins as a key macromolecule. *Molecules*. **23**:1098(2018).
- 24 100.Solé-Bundó M, Salvadó H, Passos F, Garfí M, Ferrer I. Strategies to optimize
25 microalgae conversion to biogas: co-digestion, pretreatment and hydraulic retention
26 time. *Molecules*. **23**:2096(2018).
- 27 101.Gupta RB. *Hydrogen fuel: production, transport, and storage*. Crc Press, Taylor
28 and Francis group., USA, Florida(2008).
- 29 102.Skjånes K, Rebours C, Lindblad P. Potential for green microalgae to produce
30 hydrogen, pharmaceuticals and other high value products in a combined process.
31 *Critical reviews in biotechnology*. **33**:172-215(2013).
- 32 103.Sherif SA, Goswami DY, Stefanakos EL, Steinfeld A. *Handbook of Hydrogen*
33 *Energy*. USA, Florida: CRC Press, Taylor & Francis Group; (2014).
- 34 104.Dutta D, De D, Chaudhuri S, Bhattacharya SK. Hydrogen production by
35 Cyanobacteria. *Microbial Cell Factories*. **4**:36(2005).
- 36 105.Madamwar D, Garg N, Shah V. Cyanobacterial hydrogen production. *World*
37 *Journal of Microbiology and Biotechnology*. **16**:757-67(2000).
- 38 106.Azbar N. Fundamentals of hydrogen production via biotechnology (Bio-H₂).
39 *Phytoremediation for Green Energy*: Springer., Netherlands, pp. 149-73 (2015).
- 40 107.Azwar MY, Hussain MA, Abdul-Wahab AK. Development of biohydrogen
41 production by photobiological, fermentation and electrochemical processes: A review.
42 *Renewable and Sustainable Energy Reviews*. **31**:158-73(2014).
- 43 108.Kaushik A, Sharma M. Exploiting biohydrogen pathways of cyanobacteria and
44 green algae: an industrial production approach. In: Singh A, Rathore D, editors.
45 *Biohydrogen Production: Sustainability of Current Technology and Future Perspective*.
46 Springer., New Delhi, pp. 97-113 (2017).
- 47 109.Benemann J. Hydrogen biotechnology: progress and prospects. *Nature*
48 *biotechnology*. **14**:1101(1996).

- 1 110.Meher Kotay S, Das D. Biohydrogen as a renewable energy resource-prospects and
2 potentials. *International Journal of Hydrogen Energy*. **33**:258-63(2008).
- 3 111.Nath K, Das D. Hydrogen from biomass. *Current Science*. **85**:265-71(2003).
- 4 112.Tamagnini P, Axelsson R, Lindberg P, Oxelfelt F, Wünschiers R, Lindblad P.
5 Hydrogenases and hydrogen metabolism of cyanobacteria. *Microbiology and molecular
6 biology reviews*. **66**:1-20(2002).
- 7 113.Kiss É, Mituletu M, Vass I. Transcriptional regulation of the bidirectional
8 hydrogenase by oxygen and light in two *Anabaena* species. *New Zealand Journal of
9 Botany*. **52**:28-35(2014).
- 10 114.Boison G, Bothe H, Schmitz O. Transcriptional analysis of hydrogenase genes in
11 the cyanobacteria *Anacystis nidulans* and *Anabaena variabilis* monitored by RT-PCR.
12 *Current microbiology*. **40**:315-21(2000).
- 13 115.Serebryakova LT, Medina M, Zorin NA, Gogotov IN, Cammack R. Reversible
14 hydrogenase of *Anabaena variabilis* ATCC 29413: catalytic properties and
15 characterization of redox centres. *FEBS Letters*. **383**:79-82(1996).
- 16 116.Philip D. Weyman, Brenda Pratte, Thiel aT. Transcription of *hupSL* in *Anabaena
17 variabilis* ATCC 29413 is regulated by *NtcA* and not by hydrogen *Applied and
18 environmental microbiology*. **74**:7:2103-10(2008).
- 19 117.Das D, Veziroglu TN. Advances in biological hydrogen production processes.
20 *International Journal of Hydrogen Energy*. **33**:6046-57(2008).
- 21 118.Salleh SF, Kamaruddin A, Uzir MH, Mohamed AR. Effects of cell density, carbon
22 dioxide and molybdenum concentration on biohydrogen production by *Anabaena
23 variabilis* ATCC 29413. *Energy Conversion and Management*. **87**:599-605(2014).
- 24 119.Kırtay E. Recent advances in production of hydrogen from biomass. *Energy
25 Conversion and Management*. **52**:1778-89(2011).
- 26 120.Brusca JS, Hale MA, Carrasco CD, Golden JW. Excision of an 11-kilobase-pair
27 DNA element from within the *nifD* gene in *Anabaena variabilis* heterocysts. *Journal of
28 bacteriology*. **171**:4138-45(1989).
- 29 121.Elhai J, Wolk CP. Developmental regulation and spatial pattern of expression of the
30 structural genes for nitrogenase in the cyanobacterium *Anabaena*. *The EMBO journal*.
31 **9**:3379(1990).
- 32 122.Schrautemeier B, Neveling U, Schmitz S. Distinct and differently regulated
33 Mo-dependent nitrogen-fixing systems evolved for heterocysts and vegetative cells of
34 *Anabaena variabilis* ATCC 29413: characterization of the *fdxH1/2* gene regions as part
35 of the *nif1/2* gene clusters. *Molecular microbiology*. **18**:357-69(1995).
- 36 123.Lyons EM, Thiel T. Characterization of *nifB*, *nifS*, and *nifU* genes in the
37 cyanobacterium *Anabaena variabilis*: *NifB* is required for the vanadium-dependent
38 nitrogenase. *Journal of bacteriology*. **177**:1570-5(1995).
- 39 124.Thiel T, Lyons EM, Erker JC. Characterization of genes for a second Mo-
40 dependent nitrogenase in the cyanobacterium *Anabaena variabilis*. *Journal of
41 bacteriology*. **179**:5222-5(1997).
- 42 125.Thiel T. Isolation and characterization of the *VnfEN* genes of the cyanobacterium
43 *Anabaena variabilis*. *Journal of bacteriology*. **178**:4493-9(1996).
- 44 126.Kentemich T, Haverkamp G, Bothe H. The expression of a third nitrogenase in the
45 cyanobacterium *Anabaena variabilis*. *Zeitschrift für Naturforschung C*. **46**:217-
46 22(1991).
- 47 127.Thiel T. Characterization of genes for an alternative nitrogenase in the
48 cyanobacterium *Anabaena variabilis*. *Journal of bacteriology*. **175**:6276-86(1993).

- 1 128. Ernst A, Böhme H, Böger P. Phosphorylation and nitrogenase activity in isolated
2 heterocysts from *Anabaena variabilis* (ATCC 29413). *Biochimica et Biophysica Acta*
3 (*BBA*)-*Bioenergetics*. **723**:83-90(1983).
- 4 129. Peterson RB, Wolk CP. High recovery of nitrogenase activity and of ⁵⁵Fe-labeled
5 nitrogenase in heterocysts isolated from *Anabaena variabilis*. *Proceedings of the*
6 *National Academy of Sciences*. **75**:6271-5(1978).
- 7 130. Hrouzek P, Lukešová A, Šimek M. Comparison of light and dark nitrogenase
8 activity in selected soil cyanobacteria. *Folia Microbiologica*. **49**:435-40(2004).
- 9 131. Thiel T, Pratte B. Effect on heterocyst differentiation of nitrogen fixation in
10 vegetative cells of the cyanobacterium *Anabaena variabilis* ATCC 29413. *Journal of*
11 *bacteriology*. **183**:280-6(2001).
- 12 132. Hay JXW, Wu TY, Juan JC, Md. Jahim J. Biohydrogen production through photo
13 fermentation or dark fermentation using waste as a substrate: overview, economics, and
14 future prospects of hydrogen usage. *Biofuels, Bioproducts and Biorefining*. **7**:334-
15 52(2013).
- 16 133. Bartacek J, Zabranska J, Lens PN. Developments and constraints in fermentative
17 hydrogen production. *Biofuels, Bioproducts and Biorefining: Innovation for a*
18 *sustainable economy*. **1**:201-14(2007).
- 19 134. Eroglu E, Melis A. Photobiological hydrogen production: recent advances and state
20 of the art. *Bioresource technology*. **102**:8403-13(2011).
- 21 135. Tsygankov AA, Serebryakova LT, Rao KK, Hall DO. Acetylene reduction and
22 hydrogen photoproduction by wild-type and mutant strains of *Anabaena* at different
23 CO₂ and O₂ concentrations. *FEMS Microbiology Letters*. **167**:13-7(1998).
- 24 136. Ernst A, Reich S, Böger P. Modification of dinitrogenase reductase in the
25 cyanobacterium *Anabaena variabilis* due to C starvation and ammonia. *Journal of*
26 *bacteriology*. **172**:748-55(1990).
- 27 137. Markov SA, Bazin MJ, Hall DO. Hydrogen photoproduction and carbon dioxide
28 uptake by immobilized *Anabaena variabilis* in a hollow-fiber photobioreactor. *Enzyme*
29 *and Microbial Technology*. **17**:306-10(1995).
- 30 138. Lindblad P, Christensson K, Lindberg P, Fedorov A, Pinto F, Tsygankov A.
31 Photoproduction of H₂ by wildtype *Anabaena* PCC 7120 and a hydrogen uptake
32 deficient mutant: from laboratory experiments to outdoor culture. *International Journal*
33 *of Hydrogen Energy*. **27**:1271-81(2002).
- 34 139. Ogawa RE, Carr JF. The Influence of Nitrogen on Heterocyst Production in
35 Blue-Green Algae. *Limnology and Oceanography*. **14**:342-51(1969).
- 36 140. Markov SA, Thomas AD, Bazin MJ, Hall DO. Photoproduction of hydrogen by
37 cyanobacteria under partial vacuum in batch culture or in a photobioreactor.
38 *International Journal of Hydrogen Energy*. **22**:521-4(1997).
- 39 141. Tsygankov AS, Serebryakova LT, Sveshnikov DA, Rao KK, Gogotov IN, Hall DO.
40 Hydrogen photoproduction by three different nitrogenases in whole cells of *Anabaena*
41 *variabilis* and the dependence on pH. *International Journal of Hydrogen Energy*.
42 **22**:859-67(1997).
- 43 142. Jeffries TW, Timourian H, Ward RL. Hydrogen production by *Anabaena*
44 *cylindrica*: effects of varying ammonium and ferric ions, pH, and light. *Applied and*
45 *Environmental Microbiology*. **35**:704-10(1978).
- 46 143. Jian-guo L, Hall DO, Rao KK, Tsygankov AA, Sveshnikov DA. Hydrogen
47 photoproduction of *A. variabilis* incorporated in a photobioreactor. *Chinese Journal of*
48 *Oceanology and Limnology*. **16**:118-26(1998).

144. Happe T, Schutz K, Bohme H. Transcriptional and mutational analysis of the uptake hydrogenase of the filamentous cyanobacterium *Anabaena variabilis* ATCC 29413. *Journal of bacteriology*. **182**:1624-31(2000).
145. Mikheeva Lidia E, Schmitzh O, Shestakov Sergey V, Bothe H. Mutants of the cyanobacterium *Anabaena variabilis* altered in hydrogenase activities. *Zeitschrift für Naturforschung C*. **50**:505-10(1995).
146. Sveshnikov DA, Sveshnikova NV, Rao KK, Hall DO. Hydrogen metabolism of mutant forms of *Anabaena variabilis* in continuous cultures and under nutritional stress. *FEMS Microbiology Letters*. **147**:297-301(1997).
147. Tsygankov AA, Borodin, V. B., Rao, K. K. and Hall, D. O. H₂ photoproduction by batch culture of *Anabaena variabilis* ATCC 29413 and its mutant PK84 in a photobioreactor. *Biotechnology and Bioengineering*. **64**:709–15(1999).
148. Flynn KJ, Mitra A, Greenwell HC, Sui J. Monster potential meets potential monster: pros and cons of deploying genetically modified microalgae for biofuels production. *Interface Focus*. **3**:20120037(2013).
149. Borodin VB, Tsygankov AA, Rao KK, Hall DO. Hydrogen production by *Anabaena variabilis* PK84 under simulated outdoor conditions. *Biotechnology and Bioengineering*. **69**:478-85(2000).
150. Fedorov AS, Tsygankov AA, Rao KK, Hall DO. *Anabaena variabilis* Production of hydrogen by an mutant in a photobioreactor under aerobic outdoor conditions. In: Matsunaga T, Pietro AS, editors. *Biohydrogen II*. Pergamon., Oxford, pp. 223-8 (2001).
151. Tsygankov AA, Fedorov AS, Kosourov SN, Rao KK. Hydrogen production by cyanobacteria in an automated outdoor photobioreactor under aerobic conditions. *Biotechnology and Bioengineering*. **80**:777-83(2002).
152. Sakurai H, Masukawa H, Kitashima M, Inoue K. A feasibility study of large-scale photobiological hydrogen production utilizing mariculture-raised cyanobacteria. *Recent Advances in Phototrophic Prokaryotes*: Springer., New York, pp. 291-303 (2010).
153. Mehrabadi A, Craggs R, Farid MM. Wastewater treatment high rate algal ponds (WWT HRAP) for low-cost biofuel production. *Bioresource technology*. **184**:202-14(2015).
154. Zhang Q, Yu Z, Zhu L, Ye T, Zuo J, Li X, et al. Vertical-algal-biofilm enhanced raceway pond for cost-effective wastewater treatment and value-added products production. *Water research*. **139**:144-57(2018).
155. Xin C, Addy MM, Zhao J, Cheng Y, Ma Y, Liu S, et al. Waste-to-biofuel integrated system and its comprehensive techno-economic assessment in wastewater treatment plants. *Bioresource technology*. **250**:523-31(2018).
156. Wang Y, Ho S-H, Cheng C-L, Guo W-Q, Nagarajan D, Ren N-Q, et al. Perspectives on the feasibility of using microalgae for industrial wastewater treatment. *Bioresource technology*. **222**:485-97(2016).
157. Vulsteke E, Van Den Hende S, Bourez L, Capoen H, Rousseau DP, Albrecht J. Economic feasibility of microalgal bacterial floc production for wastewater treatment and biomass valorization: A detailed up-to-date analysis of up-scaled pilot results. *Bioresource technology*. **224**:118-29(2017).
158. Gaur N, Dhankhar R. Removal of Zn⁺² ions from aqueous solution using *Anabaena variabilis*: Equilibrium and kinetic studies. *International Journal of Environmental Research*. **3**:605-16(2009).
159. Verma N, Monica, Rehal R. Biosorption of lead by alginate immobilized *Anabaena variabilis*. *Ecology, Environment and Conservation*. **2**:187-8(1996).

- 1 160.Khattar JI, Sarma TA, Singh DP, Sharma A. Bioaccumulation of chromium ions by
2 immobilized cells of a filamentous cyanobacterium, *Anabaena variabilis*. *Journal of*
3 *microbiology and biotechnology*. **12**:137-41(2002).
- 4 161.Famiglietti M, Hochkoeppler A, Luisi PL. Surfactant-Induced hydrogen production
5 in cyanobacteria. *Biotechnology and Bioengineering*. **42**:1014-8(1993).
6

Table 1. Bioremediation potential using *T. variabilis*.

Experimental conditions	Removed compounds	Efficiency (%)	Ref
Preliminary growth: MBL medium, 25–35 °C, day light for 14 days. Treatment process: Mixed domestic-industrial wastewater, 25–35 °C, daylight.	BOD	89.3	9
	COD	52.6	
	TSS	44.6	
	TDS	38.8	
	Fats, Oil and Grease	68.2	
	Zn	81.2	
	Cu	91	
Preliminary growth: BG ₁₁ medium, 30 °C for 15 days. Treatment process: ~16% of BG ₁₁ medium mixed with textile industry effluent wastewater for 25 days, 30 °C, aerobic condition.	BOD	83	10
	COD	75	
	SO ₄	55.4	
	Ni	63	
	Zn	67	
	Ca	17.5	
	Mg	28	
Preliminary growth: MDM medium, 27.5 °C, ~ 65 µmol.m ⁻² .s ⁻¹ , aerated containing 1% CO ₂ for 7 days. Treatment process: using supplemented medium, phenols concentration 40 µM, 25 °C for 120 h.	<i>o</i> -Nitrophenol	100	71
	<i>m</i> -Nitrophenol	100	
	<i>p</i> -Nitrophenol	4	
	2,4-Dinitrophenol	95	
	2,4,6-Trinitrophenol	51	
	Bisphenol A	23	
Preliminary growth: Fogg medium, 24 ± 1 °C, ~ 55µmol.m ⁻² .s ⁻¹ , 16 h light / 8 h dark cycle for 22 days. Treatment process: supplemented medium, Standard stock solution of Zn ²⁺ ions (1000 ± 2 mg.L ⁻¹)	Zn	85.1	158

Preliminary growth: calcium alginate immobilized cell Treatment process: supplemented medium, 11-10 ppm concentration of lead	Pb	96	159
Preliminary growth: Chu-10 medium, 28 ± 2 °C, daylight $\sim 45 \mu\text{mol m}^{-2} \text{s}^{-1}$ Treatment process: supplemented medium, Stock solutions of chromium concentrations of (10 - 100 mM) prepared by dissolving $\text{K}_2\text{Cr}_2\text{O}_7$.	Cr	54	160
Preliminary growth: Allen and Arnon medium, 25 °C, $15 \mu\text{mol.m}^{-2}.\text{s}^{-1}$ Treatment process: Ammonium ions were added in concentration of 0.5 mg.L^{-1} , immobilized cells in the photobioreactor for 25 days.	Ammonium	90	13

Table 2. Hydrogen production yield using *T. variabilis* under different experimental conditions.

Strain (wild/mutant)	Experimental conditions	Medium	Hydrogen production rate	Ref
ATCC 29413	Growth: 30 °C, 60 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ H ₂ : 5% CO ₂ , Ar (25 ml min ⁻¹), 35 °C, 70 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$, (a) 1.6 mM Mo, (b) 10 mM glucose	BG ₀	(a) 44 $\mu\text{mol.mg chla}^{-1}.\text{h}^{-1}$ (b) 49 $\mu\text{mol.mg chla}^{-1}.\text{h}^{-1}$	118
	Growth: 95% air + 5% CO ₂ , 30 °C, 65 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ H ₂ : Ar (45 ml min ⁻¹), 30 °C, 150 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$	BG ₀	0.9 mL.g dry cell ⁻¹ .h ⁻¹	48
	Growth: 30 °C, 35-161 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$, 190 rpm H ₂ : 100% Ar	BG ₁₁ AA BG ₁₁	1 mL.g dry cell ⁻¹ .h ⁻¹ 5.6 mL.g dry cell ⁻¹ .h ⁻¹ ~6 mL.g dry cell ⁻¹ .h ⁻¹	41
	Growth: Ambient air, 25 °C, 5 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ (bottom), 15 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ (surface) H ₂ : 100% Ar, Hydrophilic cuprammonium rayon hollow fibers PBR, medium was heated at 50°C prior injection	AA	17- 20 mg.g dry cell ⁻¹ .h ⁻¹	13
	Growth: ~ 67.5 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ H ₂ : Ar + 5% CO ₂ , 77 mM Tween 85	AA	0.44 ± 0.03 mL.mg dry cell ⁻¹ .h ⁻¹	161
CCAP 403/4B	Growth: 1.7% CO ₂ , 28 °C, continues 45-55 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$, 140 rpm H ₂ : Vacuum degassed (270-300 torr), 170-180 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ for 5 hr	AA ₀	12 .5 mL.g dry cell ⁻¹ .h ⁻¹	140
	Growth: Air without CO ₂ , 28 °C, 15 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$, 140 rpm H ₂ : No gas phase, cells immobilized on hollow fibers; 25 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ (surface), 13 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ (bottom), medium was heated at 50 °C prior injection	AA ₀	20 mL.g dry cell ⁻¹ .h ⁻¹	137

PK84	Growth: Air + 2% CO ₂ (500 mL.min ⁻¹), 12 h light (36 °C)/ 12 h dark (14- 30 °C) H ₂ : (a) air, (b) air+ 2% CO ₂ , (c) 80% Ar + 20% O ₂ , (d) 100% Ar, 332 μmol.m ⁻² .s ⁻¹	AA ₀	(a) 106.0 ± 7.0 μmol.mg chl a ⁻¹ .h ⁻¹ (b) 81.0 ± 10.0 μmol.mg chl a ⁻¹ .h ⁻¹ (c) 190.0 ± 8.0 μmol.mg chl a ⁻¹ .h ⁻¹ (d) 191.0 ± 11.0 μmol.mg chl a ⁻¹ .h ⁻¹	149
	Growth: Air+ 2% CO ₂ (500 mL.min ⁻¹) H ₂ : Outdoor PBR, ~ 1.84 mol.m ⁻² .s ⁻¹	AA ₀	80 mL.h ⁻¹ .PBR v ⁻¹	150
	Growth: air + 2% CO ₂ (500 mL.min ⁻¹), 30 °C H ₂ : Outdoor PBR, sun light	AA ₀	60-140 mL.h ⁻¹ .PBR v ⁻¹	151
(a) ATCC29413 (b) PK84	Growth: air + 2% CO ₂ (0.5 L.min ⁻¹), 30 °C, 113 μmol.m ⁻² .s ⁻¹ H ₂ : 100% Ar, 30 °C, 200 μmol.m ⁻² .s ⁻¹	AA	(a) 39.4 nmol.μg chl a ⁻¹ .h ⁻¹ (b) 32.3 nmol.μg chl a ⁻¹ .h ⁻¹	135
(a) ATCC29413 (b) PK84 (c) PK17R	Growth: 25% N ₂ , 2% CO ₂ , 73% Ar (250 mL.min ⁻¹), 90 μmol.m ⁻² .s ⁻¹ H ₂ : Ar, 30 °C, 140 μmol.m ⁻² .s ⁻¹ , N/C starvation	AA ₀	(a) 1.62- 3.07 nmol.μg prot ⁻¹ .h ⁻¹ (b) 6.91-12.6 nmol.μg prot ⁻¹ .h ⁻¹ (c) 2.24-4.10 nmol.μg prot ⁻¹ .h ⁻¹	146
AVM13	Growth: Air, 1% CO ₂ , 30 °C, 100 μmol.m ⁻² .s ⁻¹ H ₂ : Washed with N-free medium	BG ₀	68 nmol.μg ⁻¹ .chl a ⁻¹ .h ⁻¹	144